

**Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 1)**

Vaccine Requirement	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	Kindergarten through 12 <sup>th</sup> Grade		Minimum Intervals Allowed Between Doses and Other Options for Proof of Immunity
		First Entry into School (Kindergarten or First Grade)	Other Grades	
DTP/DTaP/ or Tdap, Td (Diphtheria, Tetanus, Pertussis)	Three doses by 1 year of age One additional booster dose by 2 <sup>nd</sup> birthday	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday	Three or more doses of DTP/DTaP or Td; with the last dose qualifying as a booster if received on or after the 4 <sup>th</sup> birthday  For Students entering 6 <sup>th</sup> thru 12 <sup>th</sup> grades: 1 dose of Tdap	Minimum interval between series doses: 4 weeks (28 days) Between series and booster: 6 months No proof of immunity allowed
Polio	Two doses by 1 year of age. One additional dose by 2 <sup>nd</sup> birthday	Four or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday. (progressive requirement)	Three or more doses of Polio with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday. *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a booster on or after the 4 <sup>th</sup> birthday.	Minimum interval between series doses: 4 weeks (28 days) For Grade K: 6 month interval booster must be on or after 4 <sup>th</sup> birthday No proof of immunity allowed
Measles	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Measles Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of measles immunity OR Certified physician verification* of measles disease by date of illness *Cases diagnosed after 7/1/2002 must include lab evidence of infection
Rubella	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Rubella Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of rubella immunity History of disease is not acceptable proof of immunity to rubella
Mumps	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Mumps Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		Laboratory evidence of mumps immunity OR Certified physician verification of mumps disease by date of illness
<i>Haemophilus influenzae</i> type b	Refer to Hib ACIP recs for schedule for series Children 24-59 mos. without series must have one dose after 15 mos. of age	Not required after the 5 <sup>th</sup> birthday (60 months of age)		Refer to ACIP Hib vaccination schedule  No proof of immunity allowed
Invasive Pneumococcal Disease	Refer to PCV ACIP recs for schedule for series Children 24-59 mos. without series must have one dose after 24 months of age	Not required after the 5 <sup>th</sup> birthday (60 months of age)		Refer to ACIP PCV vaccination schedule  No proof of immunity allowed

**Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 2)**

Vaccine Requirement	Kindergarten through 12 <sup>th</sup> Grade		Minimum Intervals Allowed Between Doses and Other Options for Proof of Immunity
	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	First Entry into School (Kindergarten or First Grade)	
Hepatitis B	Three doses for all children  Third dose must have been administered on or after 6 months of age (168 days)	No Requirements	Minimum intervals between doses: 1 & 2- at least 4 weeks (28 days) 2 & 3 - at least 2 months (56 days) 1 & 3 - at least 4 months (112 days) <u>Laboratory evidence of prior or current infection</u>
Varicella (progressive requirement)	One dose on or after 1 <sup>st</sup> birthday	Two doses of Varicella Vaccine, for Students entering Kindergarten and grades and 1 The 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.	Minimum intervals for administration: The 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later. <u>Statement from physician or health care provider verifying disease history OR</u> <u>Laboratory evidence of varicella immunity</u>
		Two doses of Varicella Vaccine for Students entering grades 2, 6, 7, 8, 9, 10, & 11	
Meningococcal Conjugate Vaccine (progressive requirement)	No Requirements	No Requirements	Minimum intervals for administration: The first dose received on or after the 10 <sup>th</sup> birthday; second dose at least eight weeks after the 1 <sup>st</sup> dose. Only one dose is required if the first dose was received at 16 years of age or older. <u>No proof of immunity allowed.</u>

Source: Child Health Examination Code/Part 665,  
Prepared by: Illinois Department of Public Health Immunization Section, January 2016